





| | Flu Influenza | Tdap Tetanus, diphtheria, pertussis | HPV Human papillomavirus | Meningococcal | | Pneumococcal | Dengue | Hepatitis B | Hepatitis A | Polio | MMR Measles, mumps, rubella | Chickenpox Varicella |
|--------------------------|---|--|--|--|--|--------------|--|-------------|-------------|--------|--------------------------------------|-------------------------|
| | | | | MenACWY | MenB | | | | | | | |
| 7-8 Years | Green | Orange | | Purple | | Purple | | Orange | Orange | Orange | Orange | Orange |
| 9-10 Years | Green | Orange | Partial Purple, Partial Yellow Checkered | Purple | Partial Purple | Purple | Green: ONLY in places where dengue spreads | Orange | Orange | Orange | Orange | Orange |
| 11-12 Years | Green | Orange | Green | Green | Partial Purple | Purple | Green: ONLY in places where dengue spreads | Orange | Orange | Orange | Orange | Orange |
| 13-15 Years | Green | Orange | Orange | Orange | Partial Purple | Purple | Green: ONLY in places where dengue spreads | Orange | Orange | Orange | Orange | Orange |
| 16-18 Years | Green | Orange | Orange | Orange | Partial Purple, Partial Blue | Purple | Green: ONLY in places where dengue spreads | Orange | Orange | Orange | Orange | Orange |
| More information: | Everyone 6 months and older should get a flu vaccine every year if they do not have contraindications | All 11- through 12- year olds should get one shot of Tdap. | All 11- through 12- year olds should get a 2-shot series of HPV vaccine. A 3-shot series is needed for those with weakened immune systems and those who start the series at 15 years or older. | All 11- through 12- year olds should get one shot of meningococcal conjugate (MenACWY). A booster shot is recommended at age 16. | Ages 10 years and older at increased risk should receive a serogroup B meningococcal (MenB) vaccine. Ages 16–18 years old who are not at increased risk may be vaccinated with a MenB vaccine. | | Ages 9-16 years who live in dengue endemic areas AND have laboratory confirmation of previous dengue infection | | | | | |


COVID-19 vaccination is recommended for ages 5 years and older.
Talk to your child’s doctor or nurse about the vaccines recommended for their age.

 These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

 These shaded boxes indicate the vaccine is recommended for children with certain health or lifestyle conditions that put them at an increased risk for serious diseases. See vaccine-specific recommendations at www.cdc.gov/vaccines/hcp/acip-recs/.

 These shaded boxes indicate the vaccine **SHOULD** be given if a child is catching up on missed vaccines.

 This shaded box indicates children not at increased risk **MAY** get the vaccine if they wish after speaking to a provider.

 This shaded box indicates children not at increased risk may get the vaccine if they wish after speaking to a provider.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™



AAFP
AMERICAN ACADEMY OF FAMILY PHYSICIANS

Vaccine-Preventable Diseases and the Vaccines that Prevent Them

| Disease | Vaccine | Disease spread by | Disease symptoms | Disease complications |
|------------------------------|--|--|---|---|
| Chickenpox | Varicella vaccine protects against chickenpox. | Air, direct contact | Rash, tiredness, headache, fever | Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs), death |
| Dengue | Dengue vaccine protects against dengue. | Bite from infected mosquito | May be no symptom, fever, headache, pain behind the eyes, rash, joint pain, body ache, nausea, loss of appetite feeling tired, abdominal pain | Severe bleeding, seizures, shock, damage to liver, heart, and lungs, death |
| Diphtheria | Tdap* and Td** vaccines protect against diphtheria. | Air, direct contact | Sore throat, mild fever, weakness, swollen glands in neck | Swelling of the heart muscle, heart failure, coma, paralysis, death |
| Hepatitis A | HepA vaccine protects against hepatitis A. | Direct contact, contaminated food or water | May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine | Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders, death |
| Hepatitis B | HepB vaccine protects against hepatitis B. | Contact with blood or body fluids | May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain | Chronic liver infection, liver failure, liver cancer, death |
| Human Papillomavirus | HPV vaccine protects against human papillomavirus. | Direct skin contact | May be no symptoms, genital warts | Cervical, vaginal, vulvar, penile, anal, oropharyngeal cancers |
| Influenza (Flu) | Flu vaccine protects against influenza. | Air, direct contact | Fever, muscle pain, sore throat, cough, extreme fatigue | Pneumonia (infection in the lungs), bronchitis, sinus infections, ear infections, death |
| Measles | MMR*** vaccine protects against measles. | Air, direct contact | Rash, fever, cough, runny nose, pink eye | Encephalitis (brain swelling), pneumonia (infection in the lungs), death |
| Meningococcal Disease | MenACWY and MenB vaccines protect against meningococcal disease. | Air, direct contact | Sudden onset of fever, headache, and stiff neck, dark purple rash | Loss of limb, deafness, nervous system disorders, developmental disabilities, seizure disorder, stroke, death |
| Mumps | MMR*** vaccine protects against mumps. | Air, direct contact | Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain | Meningitis (infection of the covering around the brain and spinal cord), encephalitis (brain swelling), inflammation of testicles or ovaries, deafness, death |
| Pertussis | Tdap* vaccine protects against pertussis. | Air, direct contact | Severe cough, runny nose, apnea (a pause in breathing in infants) | Pneumonia (infection in the lungs), death |
| Pneumococcal Disease | Pneumococcal vaccine protects against pneumococcal disease. | Air, direct contact | May be no symptoms, pneumonia (infection in the lungs) | Bacteremia (blood infection), meningitis (infection of the covering around the brain and spinal cord), death |
| Polio | Polio vaccine protects against polio. | Air, direct contact, through the mouth | May be no symptoms, sore throat, fever, nausea, headache | Paralysis, death |
| Rubella | MMR*** vaccine protects against rubella. | Air, direct contact | Sometimes rash, fever, swollen lymph nodes | Very serious in pregnant women—can lead to miscarriage, stillbirth, premature delivery, birth defects |
| Tetanus | Tdap* and Td** vaccines protect against tetanus. | Exposure through cuts on skin | Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever | Broken bones, breathing difficulty, death |

*Tdap combines protection against diphtheria, tetanus, and pertussis.

**Td combines protection against diphtheria and tetanus.

***MMR combines protection against measles, mumps, and rubella.

If you have any questions about your child's vaccines, talk to your child's doctor or nurse.